

# **HIGH ALTITUDE MEDICINE & BIOLOGY**

High Altitude Medicine & Biology: <http://mc.manuscriptcentral.com/highaltitudemedicine>

## **Natural Climbers: Insights from Avian Physiology at High Altitude**

Journal:	<i>High Altitude Medicine &amp; Biology</i>
Manuscript ID	HAM-2019-0032.R2
Manuscript Type:	Brief Reports
Date Submitted by the Author:	n/a
Complete List of Authors:	Parr, Nicole; University of Exeter, ; Hawkes, Lucy; University of Exeter College of Life and Environmental Sciences Wilkes, Matt; University College London, Centre for Altitude Space and Extreme Environment Medicine
Keyword:	Physiology of high altitude animals, hypobaric hypoxia, Acclimatization
Manuscript Keywords (Search Terms):	Physiology, hypoxia, comparative, birds, acclimatisation

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Natural Climbers: Insights from Avian Physiology at High Altitude**

Parr, Nicole<sup>1</sup>, Wilkes, Matt<sup>2</sup> & Hawkes, Lucy Alice<sup>3\*</sup>

1. University of Exeter, Penryn Campus, Penryn, Cornwall, TR10 9FE,  
[N.Parr@exeter.ac.uk](mailto:N.Parr@exeter.ac.uk), +44 (0) 7999 838442
2. Centre for Altitude Space and Extreme Environment Medicine, Institute of  
Sport, Exercise and Health, 170 Tottenham Court Road, London W1T 7HA,  
[M.Wilkes@ucl.ac.uk](mailto:M.Wilkes@ucl.ac.uk), +44 (0) 7976962609
3. University of Exeter, College of Life and Environmental Sciences, Hatherly  
Laboratories, Exeter EX4 4PS, UK, [L.Hawkes@exeter.ac.uk](mailto:L.Hawkes@exeter.ac.uk), +44 (0) 1392  
722248

\* Corresponding author: [L.Hawkes@exeter.ac.uk](mailto:L.Hawkes@exeter.ac.uk)

Running Title: **Avian Physiology at Altitude**

**ABSTRACT**

High altitudes are physiologically challenging: the hypobaria, hypoxia, cold, and increased UV radiation means humans ascending to high-altitude faster than they acclimatise risk life-threatening illnesses. Despite such challenges, birds can thrive at high altitudes and some even complete metabolically costly migrations across the world's highest mountain ranges. We outline the aspects of avian anatomy and physiology that confer advantages at each level of the oxygen transport cascade and compare them with those of human and non-human mammals. We also discuss additional adaptations that have been described for high-altitude specialist species of birds and how these are mirrored in high-altitude adapted mammals.

**KEYWORDS**

Physiology, hypoxia, comparative, birds, acclimatisation

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**INTRODUCTION**

High-altitude environments present a range of challenges to life, including cold, dehydration, increased exposure to solar radiation, malnutrition and hypobaric hypoxia (Bouverot, 1985). Above 2,000 m altitude humans may begin to suffer from high-altitude illnesses (reviewed in Bärtsch and Swenson, 2013). Despite this, elite human mountaineers may spend years (and tens of thousands of dollars) preparing for ascents of the world’s highest peaks (Fig. 1).

To survive in high-altitude environments, animals require a suite of physiological adjustments along the oxygen transport cascade to maintain an adequate supply of oxygen to respiring organs and tissues (Fig. 2). These adjustments may take any, or all, of three forms. First, those that are fixed as the result of generations of adaptation (the process of selection pressures increasing the expression of genes that alter physical or behavioural features at a species or population level (Willmer et al., 2005)). For example some human populations have adapted to life at high altitude, and these adaptations vary with respect to geographical origin (e.g. Tibetan versus Andean people, see reviews elsewhere (Azad et al., 2017; Beall, 2007)). Second, short-term phenotypically plastic responses (referred to as “acute hypoxia responses” from now on), which are almost instantaneous responses to hypoxia exposure, but are usually temporary and reversible (Piersma and van Gils, 2011). Third, longer-term plastic responses following continued exposure, also known as ‘acclimatisation’, which can last for days to years. Acclimatisation enables humans native to low altitudes to transiently visit extremely high altitudes, for example to climb the world’s 14 mountains over 8,000 m without the use of supplementary oxygen (Hawley et al., 2018; Wolfel et al., 1991).

Birds, regardless of the altitude at which they live, display essential features of their anatomy and physiology that put them at an advantage compared to mammals in hypoxic environments at high altitude. This was demonstrated by Tucker (1968) using a very simple hypoxic chamber experiment. After rapid ascent to a simulated altitude of 6,100 m, a mouse was rendered comatose whilst a house sparrow was able to maintain flight in a wind tunnel. The features of avian physiology that enabled this high tolerance for hypoxia are not necessarily the direct result of evolutionary adaptations following chronic hypoxic selection pressures, rather they likely evolved due to the high oxygen demands required to sustain flight. Flight is a very costly form of locomotion in terms of oxygen demand, requiring some 10-12 times resting metabolic rate (Bishop et al., 2002). Powered flight also evolved in bats, which (like birds) are also capable of high rates of oxygen consumption ( $\dot{V}O_2$ ) (reviewed in Maina, 2000). High  $\dot{V}O_2$  may therefore serve as a preadaptation for tolerating hypoxia in birds, and to some degree bats (Thomas et al., 1995), which weather radars have estimated to transiently visit altitudes up to 3,000 m (Peurach, 2001; Williams et al., 1973).

In contrast to the majority of avian species, some species of birds spend a portion of their lives at high altitude (either migrating or residing there, Fig.1) and can thus tolerate hypoxia better, due to additional adaptations or greater ability to make further temporary physiological adjustments (such species will be referred to as 'high-altitude adapted' birds from now on). The best studied high-altitude adapted avian species is the bar-headed goose (*Anser indicus*), which completes a trans-Himalayan migration, including a climb of 4,000 to 5,000 m from sea level in just 7 to 8 hours between wintering grounds in south Asia to breeding grounds in central China and Mongolia (Hawkes et al., 2011, 2012). On exposure to progressive hypoxia in laboratory studies (so removing the other stressors present at high altitude and the added costs of flight),

bar-headed geese were able to remain standing and alert at a simulated altitude of 12,190 m, whereas lowland Canada geese were unable to stand at altitudes higher than 10,668 m (Black and Tenney, 1980a), though both altitudes were considerably higher than low-altitude mammals could tolerate (Tucker, 1968).

In addition to bar-headed geese, other avian migrants fly at very high altitudes (> 5,000 m) during their flights, potentially in search of favourable conditions (Senner et al., 2018). However, to date, the highest altitudes directly recorded using tracking tags are 7,290 m for a bar-headed goose, and 6,800 m for a ruddy shelduck, although both species remain below 5,500 m for the overwhelming majority (95%) of their migration (Hawkes et al., 2011; Parr et al., 2017). As well as visitors to high altitude, several avian species that are resident at high altitude have become the subject of recent studies of high-altitude physiology and comparative transcriptomics. These species include passerines (perching birds) of the Himalayas (Grewal, 2017; Hao et al., 2019) and multiple species of waterfowl in the Andes, such as torrent duck (*Merganetta armata*, found up to 4,500 m in riverain habitats of the Andes (Dawson et al., 2016)) and Andean geese (*Neochen melanoptera*, resident at 6,000 m (Maina et al., 2017), Fig. 1). Several mammalian species also reside at high altitude, including yak (*Bos grunniens*), llama (*Lama glama*) and vicuna (*Vicugna vicugna*). The high-altitude physiology of deer mice (*Peromyscus maniculatus*) has also been increasingly studied given the large altitudinal ranges and possibility for translocation experiments between low and high altitude populations (reviewed in Storz et al., 2019). The present review contrasts the mammalian and in particular human responses to high-altitude hypoxia with those of birds, at each level of the oxygen transport cascade, highlighting where the high-altitude environment has resulted in similar and divergent phenotypic adaptations.

## 103 VENTILATION

104 In humans, the partial pressure of arterial oxygen ( $P_aO_2$ ) falls within minutes of  
105 exposure to hypobaric hypoxia. Ventilation increases in response (Fig. 2A) and, while  
106 it may temporarily arrest the decline in  $P_aO_2$ , it also causes hypocapnia; a reduction in  
107 the partial pressure of arterial carbon dioxide ( $P_aCO_2$ ), leading to respiratory alkalosis.  
108 Ultimately, this limits the initial hypoxic ventilatory response (HVR) (reviewed in  
109 Dempsey et al., 1974). Only over a period of days (duration dependent on altitude,  
110 ascent rate and individual capacity) can ventilation rates slowly rise again in  
111 lowlanders visiting high-altitude (reviewed in Hoiland et al., 2018). This is achieved by  
112 balancing respiratory alkalosis with (primarily) an increase in renal bicarbonate loss,  
113 offsetting the alkalaemia, and allowing a progressive increase in ventilation and  $P_aO_2$ .  
114 Humans resident at high altitude generally have an HVR that is similar to that of  
115 lowlanders, although a blunting of HVR has been described in Tibetans living above  
116 4,000 m (Curran et al., 1995; but for review see Gilbert-Kawai et al., 2014). High-  
117 altitude non-human mammals such as high-altitude deer mice and plateau pika  
118 (*Ochotona curzoniae*) are also able to increase total ventilation in response to acute  
119 hypoxia to a greater extent than low altitude populations (Pichon et al., 2009; reviewed  
120 in Ivy and Scott, 2017) suggesting an evolved increase in capacity for acute hypoxia  
121 responses (reviewed in Storz et al., 2010).

122 Birds too display a hypoxic ventilatory response, increasing breathing rates as  $P_aO_2$   
123 drops (Fig. 2A) (for review see Ivy and Scott, 2014). However, it takes much longer for  
124  $P_aO_2$  to fall in birds because of the remarkable avian respiratory system. In contrast to  
125 the mammalian lung, the avian lung is a unidirectionally ventilated structure, with a  
126 system of avascular air sacs distributed throughout the body that mechanically

127 ventilate a relatively rigid lung (Scheid, 1979) (Fig. 2A). The consequence of this  
128 arrangement is a lung surface that is exposed to fresh oxygenated air on both  
129 inhalation when a body of air enters the lung in passage to the anterior air sacs, then  
130 again on exhalation, as a separate body of air is expelled from the posterior air sacs  
131 through the lung (Banzett et al., 1987, 1991). Thus the avian pulmonary gas exchange  
132 surface has twice the opportunity for oxygen extraction per unit of time as the  
133 mammalian lung (Scheid et al., 1972). In addition, when  $P_aO_2$  drops low enough to  
134 stimulate a hypoxic ventilatory response, birds are able to continue hyperventilating  
135 despite hypocapnia either due to a greater capacity to buffer blood pH following  
136 respiratory alkalosis (Dodd et al., 2007; reviewed in Scott, 2011) or through  
137 insensitivity of avian brain tissue to hypocapnia (discussed below, Fig. 2D). As a result,  
138 ventilation rates are not suppressed to the same extent as in mammals.

139 In high-altitude adapted avian species, the HVR is more powerful than in lowland birds  
140 (Scott and Milsom, 2007). Bar-headed geese increase total ventilation on exposure to  
141 hypoxia by increasing tidal volume in addition to respiratory frequency, a strategy that  
142 differs from low altitude waterfowl that increase only ventilation rate (Scott and Milsom,  
143 2007). Increasing tidal volume reduces the proportional contribution of anatomical  
144 dead space ventilation, and as a result bar-headed geese sustain greater total  
145 ventilation for a given fraction of inspired oxygen than low altitude geese, suggesting  
146 they have evolved a greater capacity for acute responses to hypoxia (Lague et al.,  
147 2016). In addition, exposure to hypoxia during development appears to modify this  
148 response further: bar-headed geese raised at high altitude (3,200 m) can sustain a  
149 greater HVR than bar-headed geese raised at sea level (Lague et al., 2016). Both  
150 groups of bar-headed geese in this case may respond to declines in  $P_aO_2$  similarly,  
151 but the response in low altitude raised waterfowl is likely partially inhibited by



152 responses to hypocapnia (Lague et al., 2016). This suggests that bar-headed geese  
153 in the wild (breeding grounds in Mongolia and the Tibetan-Qinghai Plateau are  
154 between 2,000 – 4,500 m) may have an enhanced capacity for acute hypoxia  
155 responses to high-altitude hypoxia. This increased capacity is likely the result of  
156 reduced sensitivity to hypocapnia (rather than increased response to hypoxic signals)  
157 relative to conspecifics and so bar-headed geese have less restriction of minute  
158 ventilation (Scott and Milsom, 2007; Slessarev et al., 2010).

## 159 PULMONARY OXYGEN DIFFUSION

160 At the pulmonary interface, oxygen must diffuse into the blood at a rate determined by  
161 the pulmonary oxygen diffusion capacity. This is increased in Tibetan and Andean  
162 human populations through a larger pulmonary volume relative to lowlanders  
163 (Brutsaert, 2007). Larger lungs have also been found in mice (*Phyllotis darwini*) living  
164 at high altitudes (4,660 m) relative to sea level populations (Pearson and Pearson,  
165 1976). High-altitude adapted birds also have larger lungs relative to low altitude  
166 species (Scott et al., 2011), but (regardless of altitude exposure), birds have on  
167 average 15% larger lung surface areas than mammals of equivalent body mass  
168 (reviewed in West, 2009). The blood-gas barrier is also considerably thinner in birds  
169 than mammals (Fig. 2B). Non-volant mammalian blood gas barriers can range from  
170 0.33  $\mu\text{m}$  (Etruscan shrew, *Suncus etruscus*, Gehr et al., 1980) to 0.72  $\mu\text{m}$  (domestic  
171 pig, *Sus scrofa domesticus*, Meban, 1980), although heightened oxygen demand in  
172 bats has selected for a thinner blood gas barrier (range 0.20 - 0.28  $\mu\text{m}$  in the hairless  
173 bat (*Cheiromeles torquatus*) and lesser short nosed fruit bat (*Cynopterus brachyotis*)  
174 relative to non-volant mammals (Maina and King, 1984). In contrast, blood gas barriers  
175 of 0.1  $\mu\text{m}$  are not unusual in birds (such as waterfowl, sea birds and passerines but  
176 thickness can range from <0.1  $\mu\text{m}$  (hummingbirds) to 0.56  $\mu\text{m}$  in non-volant birds such

177 as ostriches, Dubach, 1981; Maina and West, 2005; reviewed in West et al., 2007).

178 Perhaps surprisingly, given the thinness of the avian blood gas barrier, the avian lung  
179 surface is much stronger than that of mammals. It is less susceptible to stress failures  
180 and more resistant to pressure than in the mammalian lung (Watson et al., 2007,  
181 2008).

182 In the avian lung, gas exchange is also far more effective than in mammals, not only  
183 as the result of unidirectional air flow, but also because of a cross-current gas  
184 exchange system (Fig. 2B), that theoretically allows  $P_{aO_2}$  to exceed the partial  
185 pressure of oxygen ( $pO_2$ ) of expired air (Scheid and Piiper, 1972). Air capillaries are  
186 intertwined with blood capillaries that deliver oxygenated blood to blood vessels  
187 running perpendicular to rigid parabronchioles, and therefore the diffusion difference  
188 of oxygen is not diminished across the length of the avian parabronchioles (reviewed  
189 in Scott, 2011). Indeed, the cross-current arrangement of the air and blood capillaries  
190 in birds may be so significant that Powell and Scheid (1989) estimated that a human  
191 climber on Mount Everest could climb 780 m higher if they had an avian parabronchial  
192 lung arrangement (reviewed in Maina, 2015). Although it has been little studied in wild  
193 flying birds (given the technical challenges), the avian lung is likely overall a far more  
194 effective gas exchange organ than the mammalian lung (but see Thomas et al., 1984).

195 In addition to the general avian advantage over mammals, high-altitude adapted birds  
196 may be able to increase pulmonary oxygen extraction over that of low-altitude species  
197 (McCracken et al., 2009b; Natarajan et al., 2015; York et al., 2017). For example, work  
198 in waterfowl (Andean geese and crested ducks) has suggested that extraction  
199 efficiency can increase with altitude up to 90% (recorded at a simulated altitude of  
200 11,000 m in Andean geese) but is ~45% at 12.2 kPa partial pressures of inspired  
201 oxygen (approximately 4,500 m). In contrast, bar-headed geese maintain a consistent

extraction efficiency of ~40% across all altitudes (Laguë et al., 2017). The increased capillarity of the blood-gas interface, and a proportionally larger lung surface area of Andean geese relative to bar-headed geese (despite the relatively larger lung of bar-headed geese in comparison to lowland waterfowl (Scott et al., 2011)) has been suggested as a possible mechanism for this difference in extraction capabilities (Laguë et al., 2017; Maina et al., 2017; York et al., 2017). Differences in Andean geese and bar-headed geese lung morphology and extraction effective in acute hypoxia highlight the variation in responses to high-altitude selection pressures present in waterfowl. This is potentially due to the differing nature of chronic or transient altitude exposure, or geographic origins (for review seee Laguë, 2017), as when the same selection pressures are present across species in the same geographic area a recent transcriptomic study found shifts in gene expression were similar across three high altitude species relative to closely related low altitude species (Hao et al., 2019).

#### **HAPE (High Altitude Pulmonary Oedema)**

In humans at low altitudes, when regions of the lung become hypoxic (due to uneven or poor ventilation), pulmonary vascular resistance increases via hypoxic pulmonary vasoconstriction, an adaptive vasomotor response that redistributes blood to better-ventilated lung segments (reviewed in Moudgil et al., 2005). However, this response can be maladaptive at altitudes where hypoxia is ubiquitous. As a result of increased or uneven blood flow to localised regions of the alveolar capillary bed (Maggiorini et al., 2001) the gas exchange barrier can become damaged, allowing proteins and erythrocytes to leak into the alveoli, restricting the potential for gas exchange, thus amplifying the vasoconstriction and stimulating an inflammatory response (reviewed in Swenson et al., 2002). This has been well-described in human mountaineers (reviewed in Basnyat and Murdoch, 2003) as a cause of non-cardiogenic pulmonary

oedema, known as HAPE (Bartsch et al., 2005, reviewed in Dehnert et al., 2007). In prolonged exposure, pulmonary arteries may thicken in response to chronic vasoconstriction and the resulting increase in pressure may cause right ventricular hypertrophy and failure (Tucker et al., 1975). Similarly in cattle at high altitude (above 2,100 m), hypoxic pulmonary hypertension has been linked to right heart failure (brisket disease) and is a major cause of livestock death in high-altitudes (Rhodes, 2005). In contrast, yaks (a closely related bovine) appear to be well adapted to the high-altitude environments of Tibet (found up to 4,500 m, Dolt et al., 2007; Qiu et al., 2012). Generations of evolution appear to have eliminated hypoxic vasoconstriction in this species (Anand et al., 1986) and yaks have thin-walled pulmonary vessels despite chronic altitude exposure (Durmowicz, 1993). In addition, to the authors' knowledge, large herbivores such as llamas which have been domesticated in high altitudes of South America 6,000–7,000 years ago do not develop right ventricular hypertrophy or pulmonary hypertension (Heath et al., 1974; Riek et al., 2019).

In contrast to the majority of mammals, birds are not known to suffer from HAPE, because, similar to the high altitude adapted yak, avian pulmonary vessels do not vasoconstrict in response to hypoxia, regardless of previous exposure of high altitudes (reviewed in West et al., 2007). Instead pulmonary pressure only increases in proportion to cardiac output (Black and Tenney, 1980b; Burton et al., 1968; Faraci et al., 1984a) and thus pulmonary oedema does not appear to occur in birds when exposed to high-altitude. Even if it did, the unidirectional ventilation and a structural honeycomb network of support struts of avian airways would ensure they could not become closed or blocked (Watson et al., 2007). However, in commercial poultry production in high-altitude environments, maladaptation from selective breeding of fast-growing chicks may have led to pulmonary arterial hypertension or 'ascites

syndrome' and cardiac failure (Rostami et al., 2016; Yersin et al., 1992). This is characterised by excessive accumulation of fluid in the abdominal cavity due to increased demands for cardiac output in hypoxic conditions that is exacerbated by breeding programmes for increased body size that is disproportional to heart and lung mass (Khajali and Wideman, 2016).

## **CIRCULATORY OXYGEN DELIVERY**

### *Cardiac output*

In addition to changes in ventilation, humans also respond to hypoxia by increasing cardiac output (Fig. 2C). Within 40 hours of exposure to a simulated altitude of 4,600 m heart rate increased by 34% in a study of resting lowlanders (Vogel and Harris, 1967). By comparison, high-altitude adapted humans from the Tibetan Plateau have a greater ability to increase heart rate, achieving higher maximal heart rates during exercise (for review see Gilbert-Kawai et al., 2014). Similarly, high-altitude deer mice were found to sustain higher heart rates than low altitude populations, both in high and low-altitude conditions (reviewed in Ivy and Scott, 2017). Birds, in general can sustain greater cardiac outputs; a 1 kg bird at rest has a cardiac output of approximately 290 mL.min<sup>-1</sup> compared to 166 mL.min<sup>-1</sup> for a mammal of the same mass ,in addition, birds have proportionally larger hearts than mammals (Calder, 1968; Grubb, 1983) and can more than double their cardiac output in hypoxia through a combination of increased heart rate and stroke volume (Black and Tenney, 1980a; Lague et al., 2016).

On acclimatisation to hypoxia, much like humans, birds can also increase cardiac output even in species already adapted to high-altitude; bar-headed geese that had undergone a four-week period at a simulated altitude of 5,450 m had a greater cardiac output than un-acclimatised bar-headed geese (by approximately 200 mL.min<sup>-1</sup>)

(Black and Tenney, 1980a). Similarly, bar-headed geese raised at high altitude increased cardiac output in response to diminished oxygen sooner than bar-headed geese raised at sea level (Lague et al., 2016), further demonstrating the importance of developmental phenotypic plastic responses in addition to acclimatisation for increasing the capacity to deliver oxygen through both increased ventilation and cardiac output (Monge and León-Velarde, 1991; Storz et al., 2010).

### *Blood properties*

In addition to acute hypoxia responses, humans also make haematological adjustments after days to weeks at high altitude to sustain oxygen delivery to systemic tissue, typically by increasing haematocrit and total body haemoglobin mass (Fig. 2C) and therefore increasing arterial oxygen content (reviewed in Monge and León-Velarde, 1991). This also appears to occur as a fixed, evolved response to high altitude hypoxia in humans native to the high-altitude Andes (Beall et al., 1998). The cost-benefit of this is uncertain; despite the potential increase in arterial oxygen content, the resulting increase in blood viscosity creates a greater cardiac load, which may impair microcirculation, thus ultimately decreasing oxygen delivery to peripheral tissues (for reviews see Barve et al., 2016; Monge and León-Velarde, 1991; Storz et al., 2010). At its extreme, prolonged elevation of haematocrit (secondary to central hypoventilation or increased bone marrow erythropoietin production) can be associated with chronic mountain sickness in humans (reviewed in Villafuerte and Corante, 2016) and may perhaps explain why this adaptation is not seen in humans native to Tibet (Beall et al., 1998) or other high-altitude adapted mammals such as Andean camelids (reviewed in Monge and León-Velarde, 1991).



Low altitude species of birds also increase haematocrit when experimentally exposed to simulated high altitude for prolonged periods (e.g. 3 weeks or more) (Mcgrath, 1971; Weatheres and Snyder, 1974; Weinstein et al., 1985). A strong erythropoietic response was found in pekin ducks that underwent a four-week acclimatisation to a simulated hypobaric hypoxia of 5,450 m, (Black and Tenney, 1980b), and haematocrit levels of wild citril finches (*Serinus citrinella*) increased by 6% after ascending to 2,000 m (Borras et al., 2010). In addition to hypoxia, high-altitude environments impose multiple challenges on vertebrates that may moderate blood properties (Morton, 1976). For example the increased oxygen demand for thermogenesis in the colder environment at high altitude may also play some role in inducing an erythropoietic response, indeed lower temperatures experienced by wild citril finches following ascents to 2,000 m were suggested to contribute to the increase in haematocrit levels (Borras et al., 2010). Thus, hypoxia alone may not be the sole driver for haematological changes.

In contrast to the pekin duck erythropoietic response, increases in haematocrit level were not seen in bar-headed geese when exposed to a simulated altitude of 5,450 m (Black and Tenney, 1980b). This may be because other adaptations in bar-headed geese, as well as some other bird species, sufficed to maintain oxygen delivery (Butler et al. 2010; Scott et al. 2015; reviewed in Monge and León-Velarde, 1991), thus avoiding the increased cardiac load that results from increasing haematocrit (Black and Tenney, 1980b). This also appears to be the case in Tibetan natives and other high altitude non-human mammals that have a blunted erythropoietic response to acute hypoxia, such as deer mice (Lui et al., 2015).

A range of species living at high altitude have haemoglobin with an increased affinity for oxygen (McCracken et al., 2009a, 2010; Natarajan et al., 2015). Both bar-headed

geese and Andean waterfowl have a lower  $P_{50}$  (the  $pO_2$  at which haemoglobin is 50% saturated) relative to other avian species (Black and Tenney, 1980b; Liang et al., 2001; Meir and Milsom, 2013), though the mutation that causes this higher affinity differs in bar-headed geese from that in Andean waterfowl (Leu- $\beta 55$  to Ser- $\beta 55$  in Andean geese compared with Pro- $\alpha 119$  to Ala- $\alpha 119$  in bar-headed geese (Weber et al., 1985)). Camelids and high altitude populations of deer mice have also been found to have haemoglobin with a higher affinity for oxygen than their low altitude counterparts (Storz, 2007). However, the overall increase in oxygen consumption that can be achieved through a high oxygen affinity is the result of a trade-off between oxygen loading and unloading, sensitive to the altitude experienced (Storz, 2007). A recent comparative analysis demonstrated that there was ambiguous evidence for increased haemoglobin oxygen affinity in high altitude mammals, although across avian species there was a strong positive relationship between bird species' native elevation (taken as the mid-point of each species' elevation range (Projecto-Garcia et al., 2013)) and haemoglobin oxygen affinity (Storz, 2016). This suggests that high altitude selection pressures on birds act to increase haemoglobin oxygen affinity and this contributes to high-altitude birds' ability to sustain adequate oxygen supply without incurring the costs of elevated haematocrit levels that lowland birds and mammals experience.

#### **CEREBRAL OXYGEN DELIVERY AND HACE (High Altitude Cerebral Oedema)**

Human lowlanders respond to hypoxia induced decreased  $P_aO_2$  with temporarily increased cerebral blood flow. Upon initial exposure to high-altitude, cerebral vasculature dilates and for a short period, blood flow and oxygen delivery to the brain increases to compensate for reduced  $P_aO_2$  (reviewed in Sanborn et al., 2015). However, this increase is subsequently diminished as a result of other acute hypoxia



responses. Firstly, increased ventilation leads to hypocapnia, which causes vasoconstriction of cerebral blood vessels (Fig. 2D). Secondly, other acclimatory changes such as increased haematocrit, result in greater arterial oxygen content, so that cerebral blood flow can decrease without diminishing total cerebral oxygen delivery (for review see Hoiland et al., 2018). Birds response to high-altitude hypoxia also begins with increased ventilation and a dilation of the brain vasculature so that cerebral blood flow is increased. However, an insensitivity of avian brain tissue to hypocapnia means that avian cerebral vasculature does not appear to vasoconstrict in response to the ensuing hypocapnia caused by increased ventilation, so cerebral blood flow is not diminished following prolonged hypoxic exposure (Faraci, 1991; Faraci et al., 1985) (Fig. 2D).

In humans, alterations to cerebral blood flow in response to hypoxia have been implicated in the onset of high-altitude cerebral oedema in climbers (HACE). HACE is thought to be a progression of acute mountain sickness that is marked by ataxia, headache, diminished conscious level and, ultimately, death (reviewed in Luks et al., 2017). It has been suggested that HACE may result from localised differences in cerebral blood flow, or where vasodilation and increased inflow are not matched to venous outflow, leading to microhaemorrhages, and a diverse range of symptoms (Sagoo et al., 2017). In birds, the lack of vasoconstriction in response to hypocapnia means that cerebral blood pressure should not increase despite the increase in cerebral blood flow (Grubb et al., 1977, 1978). Furthermore, avian neurones are less sensitive to hypoxia than mammalian neurons (Ludvigsen and Folkow, 2009), consequently HACE is not known to occur in birds (Scott, 2011).

## **SKELETAL MUSCLE OXYGEN DELIVERY AND USE**

At the tissue and cellular level, further adaptations and acute hypoxia responses improve oxygen utilisation in hypoxia in both mammals and birds. Human lowlanders visiting high altitude over a period of one to two months (i.e. the typical duration of an expedition in the Himalayas) undergo several muscular changes, including a reduction in cross-sectional surface area of muscle fibres (Hoppeler et al., 1990; MacDougall et al., 1991), and thus an improvement in oxygen diffusion as the capillary bed serves a smaller muscle volume (Hoppeler and Vogt, 2001). In addition, during a typical expedition to high altitude, the oxidative capacity of lowlanders' skeletal muscle also decreases. Oxidative capacity is the ability to consume oxygen and oxidise substrates, independent of the oxygen availability within the cell. One way this can be reduced is through a smaller volume density of mitochondria, which has been found to occur in human lowlanders who undergo a reduction of up to 30% in the subsarcolemmal or interfibrillar mitochondria following expeditions to high altitude (Hoppeler et al., 1990). Furthermore, several respiratory enzymes including citrate synthase, cytochrome oxidase (Howald et al., 1990), succinate dehydrogenase and hexokinase activities also decrease (Green et al., 1989; MacDougall et al., 1991). This serves to potentially reduce production of harmful reactive oxygen species (Hoppeler et al., 2003). For athletes wishing to incur some of the benefits of exposure to high altitude (i.e. increased oxygen carrying capacity of the blood) without incurring these muscular costs, a schedule of intermittent hypoxia may be required (reviewed in Dempsey and Morgan, 2015). Total oxygen consumption in human high altitude residents may be enhanced relative to that in lowlanders through increased oxygen delivery, achieved by greater capillary to muscle fibre ratios (Kayser et al., 1991) and higher rates of oxygen consumption per mitochondria. This may be the result of an enhanced

efficiency in ATP turn over achieved through coupling of ATP supply and demand pathways (Hoppeler and Vogt, 2001; Hoppeler et al., 2003).

However, birds have greater mass-specific  $\dot{V}O_2$  than mammals, and this is supported by up to twice the capillarity in locomotor muscles than mammals (Fig. 2D) (Faraci, 1991; Mathieu-Costello et al., 1992) meaning that oxygen delivery to the muscles is more effective, as diffusion distances are reduced. This advantage is increased further in bar-headed geese, which have significantly higher capillary densities and higher capillary to muscle fibre ratios at the surface of flight muscles, relative to those in low altitude waterfowl species, facilitating higher levels of oxygen delivery required during high-altitude flights (Scott et al. 2011; Scott et al. 2009). Furthermore, avian flight muscle has a higher oxidative capacity and greater ability to metabolise fat than mammals (Rosser and George, 1986; for reviews see Butler, 2016; McGuire and Guglielmo, 2009). When mammals are exercising at maximal rates, fatty acids supply approximately 15% of the fuel metabolised (Weber et al., 1996). However, in birds 90% of energy demand maybe derived from fatty acids, made possible by a heightened capacity to transport fatty acids through an increase in fatty acid transport proteins on the membranes of the muscles relative to mammals (Fig. 2E) (reviewed in Guglielmo, 2010). This appears to be an adaptation necessary to sustain the high oxygen demands of flight (Tucker, 1970), rather than a response to hypobaric hypoxia per se. It is yet to be shown if greater fatty acid metabolism is beneficial at high altitudes given the higher oxidative cost of metabolising fatty acids; oxidizing lipids becomes increasingly expensive relative to glycogen with increasing altitude (Melzer, 2011; Yap et al., 2018). Enzymes involved in the metabolism of fatty acids were not found to vary across the locomotory muscles of high and low altitude populations of torrent duck (Dawson et al., 2016). However, high altitude populations of deer mice

have been found to have greater proportions of type I muscle fibres than low altitude populations, which may increase their capacity to transport fatty acids (Lui et al., 2015). Furthermore, relative to low-altitude cattle, yaks have been shown to undergo modifications in five key genes in nutrition pathways relating to fatty acid metabolism (Qiu et al., 2012). These studies suggest that fatty acid metabolism may be selected for or enhanced in high-altitude environments in mammals but it is not yet clear if the same is true in birds, which already have a higher capacity to metabolise fatty acids than mammals (Lui et al., 2015a; for review see Storz et al., 2019).

Avian high-altitude species can further increase oxidative capacity at the cellular level. High-altitude resident populations of torrent duck have significantly increased activities of cytochrome oxidase, phosphofructokinase, pyruvate kinase and malate dehydrogenase in the locomotor muscles relative to those in ducks living at low altitude (Dawson et al., 2016). It is unknown whether these adaptations are common among other high-altitude adapted bird species (respiratory enzyme activity has not been studied in many other species to date) but other adaptations at the cellular level have been described in bar-headed geese. These include a redistribution of mitochondria closer to the capillaries, relative to those in low altitude waterfowl, which is the opposite response of low altitude humans acutely exposed to high-altitude, whom undergo a reduction in subsarcolemmal mitochondria (Hoppeler et al., 1990; Scott et al., 2009, 2011). Similar adaptations have also been described in high altitude populations of deer mice, including increased capillarity, a redistribution of mitochondria to reduce the intracellular oxygen diffusion distances and increased activity of key respiratory enzymes (Lui et al., 2015b; Mahalingsam et al., 2017; Scott et al., 2015a; for review see McClelland and Scott, 2018). Therefore, chronic high-altitude hypoxia exposure appears to have acted in both birds and mammals to enhance the oxidative capacity

as well as oxygen delivery to skeletal muscle. Such increases in oxidative capacity at the skeletal muscle contribute to greater overall maximum oxygen consumption ( $\dot{V}O_{2max}$ ). At the taxon level birds appear to be able to support a higher  $\dot{V}O_{2max}$  than mammals (Fig. 2), which is beneficial at high altitudes characterised by both reduced oxygen availability and increased thermoregulation demands given the cold challenge.

## CONCLUSION

Birds' ability to sustain the high metabolic rates to facilitate the demands of flight likely confer benefits that allow birds to better tolerate hypoxia relative to mammals, including humans. Species and populations of birds and mammals that are regularly exposed to high-altitudes have adaptations to enhance oxygen delivery, including greater HVR and a greater capacity for oxygen carrying capacity of the blood (reviewed in Butler, 2010; Ivy and Scott, 2014; Scott, 2011; Scott et al., 2015). More recent research has shown that high altitude can also act as a strong selection pressure for greater oxidative capacity at the cellular level across both birds (Dawson et al., 2016; Hao et al., 2019; Yap et al., 2018) and mammals (McClelland and Scott, 2018; Storz, 2007). Despite these shared features, the advantages of the avian lung, haemoglobin with a high oxygen affinity and insensitivity to hypocapnia triumphs. Recent comparisons of high-altitude adaptations in resident waterfowl in the Andes and migratory bar-headed geese reveal that different mechanisms can serve to increase hypoxia tolerance (Laguë et al., 2017; reviewed in Laguë, 2017). However there are common features unique to birds that visit high altitudes that include more effective gas exchange with a greater hypoxic ventilatory response, larger lungs with higher capillarization, enhanced cardiac output, greater oxygen delivery to skeletal muscle due to increased haemoglobin oxygen affinity, and higher capillary to fibre ratios with increased oxidative capacity. Indeed, on rare occasions, several species of

1  
2  
3 472 birds have been detected flying at exceptional altitudes (Hawkes et al., 2012;  
4  
5 473 Laybourne, 1974; Noel, 1927; Parr et al., 2017). While extreme high-altitude flight may  
6  
7 474 not represent their typical behaviour, the question remains as to how high the best  
8  
9 475 adapted avian species could reach, possibly with the assistance of orographic lift  
10  
11  
12 476 (Bishop et al., 2015).

13  
14  
15 477 **Authorship Confirmation Statement**

16  
17  
18 478 All authors contributed to the conception of the manuscript; research and first drafting  
19  
20 479 of the manuscript was conducted by NP. MW and LH contributed to the subsequent  
21  
22 480 writing and editing. All authors have reviewed and approved the manuscript for  
23  
24  
25 481 submission.

26  
27  
28 482 **Author Disclosure**

29  
30  
31 483 We have no affiliations to declare

32  
33  
34 484 **FIGURE LEGENDS**

35  
36  
37 485 **Figure 1:** Schematic diagram showing records of humans and birds found at high  
38  
39 486 altitude in relation to commercial aircraft cruising altitude and some of the world's  
40  
41 487 highest mountains. High-altitude birds are, for the majority of the time, found at much  
42  
43 488 lower altitudes, but make rare flights above 5,500 metres altitude. Sources: Beall,  
44  
45 489 2014; Borrás et al., 2010; Dawson et al., 2016; Hawkes et al., 2011; Hawley et al.,  
46  
47 490 2018; Laybourne, 1974; McCracken et al., 2009b; Mukherjee et al., 2008; Noel, 1927;  
48  
49  
50 491 Parr et al., 2017; West, 2002

51  
52  
53 492 **Figure 2:** Diagram of the oxygen transport chain, detailing the physiological  
54  
55 493 adjustments to high-altitude hypoxia that have been documented in low-altitude native  
56  
57 494 humans ascending to high altitude and details of preadaptations that are common  
58  
59  
60 495 across all birds and allow for a greater tolerance of hypoxia relative to mammals before

any physiological adjustments occur upon exposure to hypoxia. Sources: [1] West, 2006, [2] Moudgil et al., 2005, [3] Grocott et al., 2009, [4] Wagner, 2010, [5] Hoiland et al., 2018, [6] Maina, 2000, [7] Scott, 2011, [8] Scheid and Piiper, 1972, [9] Lague et al., 2016, [10] Grubb, 1983, [11] Maginniss et al., 1997, [12] Scott et al., 2009, [13] Faraci et al., 1984, [14] McGuire and Guglielmo, 2009, [15] Ward et al., 2002.



## 502 REFERENCES

- 503 Anand IS, Harris E, Ferrari R, Pearce P, and Harris P (1986). Pulmonary  
504 haemodynamics of the yak, cattle, and cross breeds at high altitude. *Thorax* 696–700.
- 505 Azad P, Stobdan T, Zhou D, Hartley I, Akbari A, Bafna V, and Haddad GG (2017).  
506 High-altitude adaptation in humans : from genomics to integrative physiology. *J Mol*  
507 *Med* 95:1269–1282.
- 508 Banzett RB, Butler JP, Nations CS, Barnas GM, Lehr JL, and Jones JH (1987).  
509 Inspiratory aerodynamic valving in goose lungs depends on gas density and velocity.  
510 *Respir Physiol* 70:287–300.
- 511 Banzett RB, Nations CS, Wang N, Fredberg JJ, and Butler JP (1991). Pressure  
512 profiles show features essential to aerodynamic valving in geese. *Respir Physiol*  
513 84:295–309.
- 514 Bartsch P, Mairba H, Maggiorini M, and Swenson ER (2005). Physiological aspects of  
515 high-altitude pulmonary edema. *J Appl Physiol* 98:1101–1110.
- 516 Bärtsch P and Swenson ER (2013). Acute High-Altitude Illnesses. *N Engl J Med*  
517 368:2294–2302.
- 518 Barve S, Dhondt AA, Mathur VB, and Cheviron ZA (2016). Life-history characteristics  
519 influence physiological strategies to cope with hypoxia in Himalayan birds. *Proc R Soc*  
520 *B Biol Sci* 283.
- 521 Basnyat B and Murdoch DR (2003). High-altitude illness. *Lancet* 361:1967–1974.
- 522 Beall CM (2007). Two routes to functional adaptation: Tibetan and Andean high-  
523 altitude natives. *Proc Natl Acad Sci U S A* 104:8655–8660.



- 524 Beall CM (2014). Adaptation to High Altitude : Phenotypes and Genotypes. *Annu Rev*  
525 *Anthr* 43:251–272.
- 526 Beall CM, Brittenham GM, Strohl KP, Blangero J, Williams-blangero S, Goldstein MC,  
527 Decker MJ, Vargas E, Villena M, Soria R, Alarcon ANAM, and Gonzales C (1998).  
528 Hemoglobin Concentration of High-Altitude Tibetans and Bolivian Aymara. *Am J Phys*  
529 *Anthropol* 400:385–400.
- 530 Bishop CM, Ward S, Woakes AJ, and Butler PJ (2002). The energetics of barnacle  
531 geese (*Branta leucopsis*) flying in wild and captive conditions. *Comp Biochem Physiol*  
532 133:225–237.
- 533 Bishop CM, Hawkes LA, Chua B, Frappell PB, Milsom WK, Natsagdorj T, Newman  
534 SH, Scott GR, Takekawa JY, Wikelski M, and Butler PJ (2015). The roller coaster flight  
535 strategy of bar-headed geese conserves energy during Himalayan migrations.  
536 *Science* (80- ) 347:250–254.
- 537 Black CP and Tenney SM (1980a). Oxygen transport during progressive hypoxia in  
538 high-altitude and sea-level waterfowl. *Respir Physiol* 39:217–239.
- 539 Black CP and Tenney SM (1980b). Pulmonary hemodynamic responses to acute and  
540 chronic hypoxia in two waterfowl species. *Comp Biochem Physiol* 67A:291–293.
- 541 Borrás A, Cabrera J, and Senar J. (2010). Hematocrit Variation in Response to Altitude  
542 Changes in Wild Birds: A Repeated-Measures Design. *Condor* 112:622–626.
- 543 Bouverot P (1985). General Introduction. In: *Adaptation to Altitude-Hypoxia in*  
544 *Vertebrates. Zoophysiology*. Springer: Berlin, Heidelberg, pp. 1–2.
- 545 Brutsaert TD (2007). Population genetic aspects and phenotypic plasticity of  
546 ventilatory responses in high altitude natives. *Respir Physiol Neurobiol* 158:151–160.

- 547 Burton RR, Besch EL, and Smith AH (1968). Effect of chronic hypoxia on the  
548 pulmonary arterial blood pressure of the chicken. *Am J Physiol* 214:1438–1442.
- 549 Butler PJ (2010). High fliers: The physiology of bar-headed geese. *Comp Biochem*  
550 *Physiol Part A Mol Integr Physiol* 156:325–329.
- 551 Butler PJ (2016). The physiological basis of bird flight. *Philos Trans R Soc Lond B Biol*  
552 *Sci* 371.
- 553 Calder W (1968). Respiratory and Heart Rates of Birds at Rest. *Condor* 70:358–365.
- 554 Curran LS, Zhuang J, Droma T, Land L, and Moore LG (1995). Hypoxic ventilatory  
555 responses in Tibetan residents of 4400 m compared with 3658 m. *Respir Physiol*  
556 100:223–230.
- 557 Dawson NJ, Ivy CM, Alza L, Cheek R, York JM, Chua B, Milsom WK, McCracken KG,  
558 and Scott GR (2016). Mitochondrial physiology in the skeletal and cardiac muscles is  
559 altered in torrent ducks, *Merganetta armata*, from high altitudes in the Andes. *J Exp*  
560 *Biol* 219:3719–3728.
- 561 Dehnert C, Berger MM, Mairbäurl H, and Bärtsch P (2007). High altitude pulmonary  
562 edema: A pressure-induced leak. *Respir Physiol Neurobiol* 158:266–273.
- 563 Dempsey JA and Morgan BJ (2015). Humans In Hypoxia: A Conspiracy Of  
564 Maladaptation?! *Physiology* 30:304–316.
- 565 Dempsey J., Forster H., and Dopico G. (1974). Ventilatory Acclimatization to Moderate  
566 Hypoxemia in Man. *J Clin Invest* 53:1091–1100.
- 567 Dodd GAA, Scott GR, and Milsom WK (2007). Ventilatory roll off during sustained  
568 hypercapnia is gender specific in pekin ducks. *Respir Physiol Neurobiol* 156:47–60.

- 569 Dolt KS, Mishra MK, Karar J, Baig MA, Ahmed Z, and Pasha MAQ (2007). cDNA  
570 cloning , gene organization and variant specific expression of HIF-1  $\alpha$  in high altitude  
571 yak (*Bos grunniens*). *Gene* 386:73–80.
- 572 Dubach M (1981). Quantitative analysis of the respiratory system of the house  
573 sparrow, budgerigar and violet-eared hummingbird. *Respir Physiol* 46:43–60.
- 574 Durmowicz A. (1993). Functional and structural adaptation of the yak pulmonary  
575 circulation to residence at high altitude. *J Appl Physiol* 74:2276–2285.
- 576 Faraci FM (1991). Adaptations to hypoxia in birds: how to fly high. *Annu Rev Physiol*  
577 53:59–70.
- 578 Faraci FM, Kilgore Jr. DL, and Fedde MR (1984a). Attenuated pulmonary pressor  
579 response to hypoxia in bar-headed geese. *Am J Physiol* 247:R402-3.
- 580 Faraci FM, Kilgore, D. L. J, and Fedde MR (1984b). Oxygen delivery to the heart and  
581 brain during hypoxia: Pekin duck vs. bar-headed goose. *Am J Physiol Regul Integr*  
582 *Comp Physiol* 247:R69-75.
- 583 Faraci FM, Kilgore DL, and Fedde MR (1985). Blood flow distribution during  
584 hypocapnic hypoxia in pekin ducks and bar-headed geese. 21–30.
- 585 Gehr P, Sehovic S, Burri P, Claassen H, and Weibel E (1980). The lung of shrews:  
586 morphometric esestimateion of diffusion capacity. *Respir Physiol* 40:33–47.
- 587 Gilbert-Kawai ET, Milledge JS, Grocott MPW, and Martin DS (2014). King of the  
588 Mountains: Tibetan and Sherpa Physiological Adaptations for Life at High Altitude.  
589 *Physiology* 29:388–402.
- 590 Green HJ, Young PM, Sutton JR, Cymerman A, and Houston CS (1989). Operation  
591 Everest II: adaptations in human skeletal muscle. *J Appl Physiol* 66:2454–2461.

- 592 Grewal B (2017). Birds of the Himalayas. Bloomsbury publishing.
- 593 Grocott M, Martin D, Levett D, McMorrow R, Windsor J, and Montgomery H (2009).  
594 Arterial Blood Gases and Oxygen Content in Climbers on Mount Everest. *N Engl J*  
595 *Med* 360:140–149.
- 596 Grubb BR (1983). Allometric relations of cardiovascular function in birds. *Am J Physiol*  
597 245:567–572.
- 598 Grubb B, Mills CD, Colacino JM, and Schmidt-Nielsen K (1977). Effect of arterial  
599 carbon dioxide on cerebral blood flow in ducks. *Am J Physiol Circ Physiol* 232:H596–  
600 H601.
- 601 Grubb B, Colacino JM, and Schmidt-Nielsen K (1978). Cerebral blood flow in birds:  
602 effect of hypoxia. *Am J Physiol* 234:H230–H234.
- 603 Guglielmo CG (2010). Move that fatty acid: Fuel selection and transport in migratory  
604 birds and bats. *Integr Comp Biol* 50:336–345.
- 605 Hao Y, Xiong Y, Cheng Y, Song G, Jia C, Qu Y, and Lei F (2019). Comparative  
606 transcriptomics of 3 high-altitude passerine birds and their low-altitude relatives. *Proc*  
607 *Natl Acad Sci* 1–6.
- 608 Hawkes L, Balachandran S, Batbayar N, Butler PJ, Frappell PB, Milsom WK,  
609 Tseveenmyadag N, Newman SH, Scott GR, Sathiyaselvam P, Takekawa JY, Wikelski  
610 M, and Bishop CM (2011). The trans-Himalayan flights of bar-headed geese (*Anser*  
611 *indicus*). *Proc Natl Acad Sci U S A* 108:9516–9519.
- 612 Hawkes L, Balachandran S, Batbayar N, Butler PJ, Chua B, Douglas DC, Frappell PB,  
613 Hou Y, Milsom WK, Newman SH, Prosser DJ, Sathiyaselvam P, Scott GR, Takekawa  
614 JY, Natsagdorj T, Wikelski M, Witt MJ, Yan B, and Bishop CM (2012). The paradox of

- 615 extreme high-altitude migration in bar-headed geese *Anser indicus*. *Proc R Soc B Biol*  
616 *Sci* 280.
- 617 Hawley E, Bierling B, Shrestha J, Popier R, Pantel T, Salisbury R, and Choegyal L  
618 (2018). The Himalayan Database.
- 619 Heath D, Smith P, Williams D, Harris P, Arias-Stella J, and Krüger H (1974). The heart  
620 and pulmonary vasculature of the llama (*Lama glama*). *Thorax* 29:463–471.
- 621 Hoiland RL, Howe CA, Coombs GB, and Ainslie PN (2018). Ventilatory and  
622 cerebrovascular regulation and integration at high-altitude. *Clin Auton Res* 28:1–13.
- 623 Hoppeler H and Vogt M (2001). Muscle tissue adaptations to hypoxia. *J Exp Biol*  
624 204:3133–3139.
- 625 Hoppeler H, Kleinert E, Howald H, Claassen H, Cerretelli P, Kayser S, and Schlegel C  
626 (1990). Morphological Adaptations of Human Skeletal Muscle to Chronic Hypoxia. *Int*  
627 *J Sports Med* 11:3–9.
- 628 Hoppeler H, Vogt M, Weibel RE, and Fluck M (2003). Response of skeletal muscle  
629 mitochondria to hypoxia. *Exp Physiol* 88:109–119.
- 630 Howald H, Cerretelli P, Pette D, Uber A, Simoneau J-A, and Hoppeler H (1990). Effects  
631 of Chronic Hypoxia on Muscle Enzyme Activities. *Int J Sports Med* 11:10–14.
- 632 Ivy CM and Scott GR (2014). Control of breathing and the circulation in high-altitude  
633 mammals and birds. *Comp Biochem Physiol A Mol Integr Physiol* 186:66–74.
- 634 Ivy CM and Scott GR (2017). Control of breathing and ventilatory acclimatization to  
635 hypoxia in deer mice native to high altitudes. *Acta Physiol* 221:266–282.
- 636 Kayser B, Hoppeler H, Claassen H, and Cerretelli P (1991). Muscle structure and

- performance of Himalayan Sherpas. *J Appl Physiol* 70:1938–1942.
- Khajali F and Wideman RF (2016). Nutritional approaches to ameliorate pulmonary hypertension in broiler chickens. *J Anim Physiol Anim Nutr (Berl)* 100:3–14.
- Lague SL, Chua B, Farrell AP, Wang Y, and Milsom WK (2016). Altitude matters: differences in cardiovascular and respiratory responses to hypoxia in bar-headed geese reared at high and low altitudes. *J Exp Biol* 219:1974–1984.
- Laguë SL (2017). High-altitude champions: birds that live and migrate at altitude. *J Appl Physiol* 123:942–950.
- Laguë SL, Chua B, Alza L, Scott GR, Frappell PB, Zhong Y, Farrell AP, McCracken KG, Wang Y, and Milsom WK (2017). Divergent respiratory and cardiovascular responses to hypoxia in bar-headed geese and Andean birds. *J Exp Biol* 220:4186–4194.
- Laybourne R (1974). Collision between a vulture and an aircraft at an altitude of 37,000 feet. *Wilson Bull* 86:461–462.
- Liang Y, Hua Z, Liang X, Xu Q, and Lu G (2001). The crystal structure of bar-headed goose hemoglobin in deoxy form: the allosteric mechanism of a hemoglobin species with high oxygen affinity. *J Mol Biol* 313:123–137.
- Ludvigsen S and Folkow LP (2009). Differences in vitro cerebellar neuronal responses to hypoxia in eider ducks, chicken and rats. *J Comp Physiol A Neuroethol Sensory, Neural, Behav Physiol* 195:1021–1030.
- Lui MA, Mahalingam S, Patel P, Connaty AD, Ivy CM, Cheviron ZA, Storz JF, McClelland GB, and Scott GR (2015). High-altitude ancestry and hypoxia acclimation have distinct effects on exercise capacity and muscle phenotype in deer mice. *Am J*

- Physiol - Regul Integr Comp Physiol 308:R779–R791.
- Luks AM, Swenson ER, and Bärtsch P (2017). Acute high-altitude sickness. *Eur Respir Rev* 26:1–14.
- MacDougall JD, Green HJ, Sutton JR, Coates G, Cymerman A, Young P, and Houston CS (1991). Operation Everest II: Structural adaptations in skeletal muscle in response to extreme simulated altitude. *Acta Physiol Scand* 142:421–427.
- Maggiorini M, Mélot C, Pierre S, Pfeiffer F, Greve I, Sartori C, Lepori M, Hauser M, Scherrer U, and Naeije R (2001). High-Altitude Pulmonary Edema Is Initially Caused by an Increase in Capillary Pressure. *Circulation* 103:2078–2083.
- Maginniss LA, Bernstein MH, Deitch MA, and Pinshow B (1997). Effects of chronic hypobaric hypoxia on blood oxygen binding in pigeons. *J Exp Zool* 277:293–300.
- Mahalingam S, McClelland GB, and Scott GR (2017). Evolved changes in the intracellular distribution and physiology of muscle mitochondria in high-altitude native deer mice. *J Physiol* 595:4785–4801.
- Maina JN (2000). What it takes to fly: the structural and functional respiratory refinements in birds and bats. *J Exp Biol* 203:3045–3064.
- Maina JN (2015). The design of the avian respiratory system: development, morphology and function. *J Ornithol* 156:41–63.
- Maina JN and King AS (1984). Correlations Between Structure and Function in the Design of the Bat Lung: a Morphometric Study. *J Exp Biol* 104:43–61.
- Maina JN and West JB (2005). Thin and strong! The bioengineering dilemma in the structural and functional design of the blood-gas barrier. *Physiol Rev* 85:811–844.



- 682 Maina JN, McCracken KG, Chua B, York JM, and Milsom WK (2017). Morphological  
683 and morphometric specializations of the lung of the Andean goose, *Chloephaga*  
684 *melanoptera*: A lifelong high-altitude resident. *PLoS One* 12:1–19.
- 685 Mathieu-Costello O, Suarez RK, and Hochachka PW (1992). Capillary-to-fiber  
686 geometry and mitochondrial density in hummingbird flight muscle. *Respir Physiol*  
687 89:113–132.
- 688 McClelland GB and Scott GR (2018). Evolved Mechanisms of Aerobic Performance  
689 and Hypoxia Resistance in High-Altitude Natives. *Annu Rev Physiol* 81:561–583.
- 690 McCracken KG, Barger CP, Bulgarella M, Johnson KP, Sonsthagen S a., Trucco J,  
691 Valqui TH, Wilson RE, Winker K, and Sorenson MD (2009a). Parallel evolution in the  
692 major haemoglobin genes of eight species of Andean waterfowl. *Mol Ecol* 18:3992–  
693 4005.
- 694 McCracken KG, Barger CP, Bulgarella M, Johnson KP, Kuhner MK, Moore AV, Peters  
695 JL, Trucco J, Valqui TH, Winker K, and Wilson RE (2009b). Signatures of High-Altitude  
696 Adaptation in the Major Hemoglobin of Five Species of Andean Dabbling Ducks. *Am*  
697 *Nat* 174:631–650.
- 698 McCracken KG, Barger CP, and Sorenson MD (2010). Phylogenetic and structural  
699 analysis of the HbA and HbD hemoglobin genes in two high-altitude waterfowl from  
700 the Himalayas and the Andes: Bar-headed goose (*Anser indicus*) and Andean goose  
701 (*Chloephaga melanoptera*). *Mol Phylogenet Evol* 56:649–658.
- 702 McGrath J (1971). Acclimation response of pigeons to simulated high altitude. *J Appl*  
703 *Physiol* 31:23–25.
- 704 McGuire LP and Guglielmo CG (2009). What Can Birds Tell Us about the Migration



- Physiology of Bats? *J Mammal* 90:1290–1297.
- Meban C (1980). Thickness of the air-blood barriers in vertebrate lungs. *J Anat* 131:299–307.
- Meir JU and Milsom WK (2013). High thermal sensitivity of blood enhances oxygen delivery in the high-flying bar-headed goose. *J Exp Biol* 216:2172–2175.
- Melzer K (2011). Carbohydrate and fat utilization during rest and physical activity. *Eur J Clin Nutr Metab* 6:e45–e52.
- Monge C and León-Velarde F (1991). Physiological adaptation to high altitude: oxygen transport in mammals and birds. *Physiol Rev* 71:1135–1172.
- Morton ML (1976). Aspects of circulatory physiology of montane and lowland birds. *Comp Biochem Physiol* 54.
- Moudgil R, Michelakis ED, Archer SL, Michelakis ED, and Hypoxic SLA (2005). Hypoxic pulmonary vasoconstriction. *Pulm Circ Hypoxia* 390–403.
- Mukherjee A, Borad C, and Parasharya B (2008). White-winged Diuca Finch (*Diuca speculifera*) Nesting on Quelccaya Ice cap, Peru. *Wilson J Ornithol* 120:89–94.
- Natarajan C, Projecto-Garcia J, Moriyama H, Weber RE, Munoz-Fuentes V, Green AJ, Kopuchian C, Tubaro PL, Alza L, Bulgarella M, Smith MM, Wilson RE, Fago A, McCracken KG, and Storz JF (2015). Convergent Evolution of Hemoglobin Function in High-Altitude Andean Waterfowl Involves Limited Parallelism at the Molecular Sequence Level. *PLoS Genet* 11:1–25.
- Noel J (1927). *Through Tibet to Everest*. E. Arnold & Company.
- Parr N, Bearhop S, Douglas DC, Takekawa JY, Prosser DJ, Newman SH, Perry WM,

- 727 Balachandran S, Witt MJ, Hou Y, Luo Z, and Hawkes LA (2017). High altitude flights  
728 by ruddy shelduck *Tadorna ferruginea* during trans-Himalayan migrations. *J Avian Biol*  
729 48:1–6.
- 730 Pearson OP and Pearson AK (1976). A stereological analysis of the ultrastructure of  
731 the lungs of wild mice living at low and high altitude. *J Morphol* 150:359–368.
- 732 Peurach SC (2001). High-altitude Collision between an Airplane and a Hoary Bat,  
733 *Lasiurus cinereus*. *Bat Res News* 44:2–3.
- 734 Pichon A, Zhenzhong B, Favret F, Jin G, Shufeng H, Marchant D, Richalet J-P, and  
735 Ge R-L (2009). Long-term ventilatory adaptation and ventilatory response to hypoxia  
736 in plateau pika ( *Ochotona curzoniae* ): role of nNOS and dopamine . *Am J Physiol*  
737 *Integr Comp Physiol* 297:R978–R987.
- 738 Piersma T and van Gils JA (2011). *The Flexible Phenotype*. Oxford University Press:  
739 Oxford.
- 740 Powell F and Scheid P (1989). Physiology of gas exchange in the avian respiratory  
741 system. *Form Funct Birds*.
- 742 Projecto-Garcia J, Natarajan C, Moriyama H, Weber RE, Fago A, Cheviron ZA, Dudley  
743 R, McGuire JA, Witt CC, and Storz JF (2013). Repeated elevational transitions in  
744 hemoglobin function during the evolution of Andean hummingbirds. *Proc Natl Acad*  
745 *Sci U S A* 110:20669–20674.
- 746 Qiu Q, Zhang G, Ma T, Qian W, Wang J, Ye Z, Cao C, Hu Q, Kim J, Larkin DM, Auvil  
747 L, Capitanu B, Ma J, Lewin HA, Qian X, Lang Y, Zhou R, Wang L, Wang K, Xia J, Liao  
748 S, Pan S, Lu X, Hou H, Wang Y, Zang X, Yin Y, Ma H, Zhang J, Wang Z, Zhang Y,  
749 Zhang D, Yonezawa T, Hasegawa M, Zhong Y, Liu W, Zhang Y, Huang Z, Zhang S,

- Long R, Yang H, Wang J, Lenstra JA, Cooper DN, Wu Y, Wang J, Shi P, Wang J, and Liu J (2012). The yak genome and adaptation to life at high altitude. *Nat Genet* 44:6–11.
- Rhodes J (2005). Comparative physiology of hypoxic pulmonary hypertension: historical clues from brisket disease. *J Appl Physiol* 98:1092–1100.
- Riek A, Stölzl A, Marquina Bernedo R, Ruf T, Arnold W, Hambly C, Speakman JR, and Gerken M (2019). Energy expenditure and body temperature variations in llamas living in the High Andes of Peru. *Sci Rep* 9:1–11.
- Rosser BWC and George JC (1986). The avian pectoralis: histochemical characterisation and distribution of muscle fiber types. *Can J Zool* 64:1174–1185.
- Rostami A, Moghaddam AKZ, Hassanpour H, and Khajali F (2016). Pulmonary hypertension and right ventricular failure in broiler chickens reared at high altitude is affected by dietary source of n-6 and n-3 fatty acids. *J Anim Physiol Anim Nutr (Berl)* 100:701–706.
- Sagoo RS, Hutchinson CE, Wright A, Handford C, Parsons H, Sherwood V, Wayte S, Nagaraja S, Ng'Andwe E, Wilson MH, and Imray CH (2017). Magnetic Resonance investigation into the mechanisms involved in the development of high-altitude cerebral edema. *J Cereb Blood Flow Metab* 37:319–331.
- Sanborn MR, Edsell ME, Kim MN, Mesquita R, Putt ME, Imray C, Yow H, Wilson MH, Yodh AG, Grocott M, and Martin DS (2015). Cerebral Hemodynamics at Altitude: Effects of Hyperventilation and Acclimatization on Cerebral Blood Flow and Oxygenation. *Wilderness Environ Med* 26:133–141.
- Scheid P (1979). Mechanisms of gas exchange in bird lungs. Springer, Berlin,

Heidelberg.

Scheid P and Piiper J (1972). Cross-current gas exchange in avian lungs: Effects of reversed parabronchial air flow in ducks. *Respir Physiol* 16:304–312.

Scheid P, Slama H, and Piiper J (1972). Mechanisms of Unidirectional Flow in Parabronchi. *Respiration* 83–95.

Scott GR (2011). Elevated performance: the unique physiology of birds that fly at high altitudes. *J Exp Biol* 214:2455–2462.

Scott GR and Milsom WK (2007). Control of breathing and adaptation to high altitude in the bar-headed goose. *Am J Physiol Regul Integr Comp Physiol* 293:379–391.

Scott GR, Egginton S, Richards JG, and Milsom WK (2009). Evolution of muscle phenotype for extreme high altitude flight in the bar-headed goose. *Proc Biol Sci* 276:3645–3653.

Scott GR, Schulte PM, Egginton S, Scott ALM, Richards JG, and Milsom WK (2011). Molecular evolution of cytochrome c oxidase underlies high-altitude adaptation in the bar-headed goose. *Mol Biol Evol* 28:351–363.

Scott GR, Elogio TS, Lui MA, Storz JF, and Cheviron ZA (2015a). Adaptive Modifications of Muscle Phenotype in High-Altitude Deer Mice Are Associated with Evolved Changes in Gene Regulation. *Mol Biol Evol* 32:1962–1976.

Scott GR, Hawkes LA, Frappell PB, Butler PJ, Bishop CM, and Milsom WK (2015b). How Bar-Headed Geese Fly Over the Himalayas. *Physiology* 30:107–115.

Senner NR, Stager M, Verhoeven MA, Cheviron ZA, Piersma T, and Bouten W (2018). High-altitude shorebird migration in the absence of topographical barriers: avoiding high air temperatures rather than searching for profitable winds. *Proc R Soc Lond B*

796 285:20180569.

797 Slessarev M, Prisman E, Ito S, Watson RR, Jensen D, Preiss D, Greene R, Norboo T,  
798 Stobdan T, Diskit D, Norboo A, Kunzang M, Appenzeller O, Duffin J, and Fisher JA  
799 (2010). Differences in the control of breathing between Himalayan and sea-level  
800 residents. *J Physiol* 588:1591–1606.

801 Storz JF (2007). Hemoglobin Function and Physiological Adaptation to Hypoxia in  
802 High-Altitude Mammals. *J Mammal* 88:24–31.

803 Storz JF (2016). Hemoglobin-oxygen affinity in high-altitude vertebrates: is there  
804 evidence for an adaptive trend? *J Exp Biol* 219:3190–3203.

805 Storz JF, Scott GR, and Cheviron ZA (2010). Phenotypic plasticity and genetic  
806 adaptation to high-altitude hypoxia in vertebrates. *J Exp Biol* 213:4125–4136.

807 Storz JF, Cheviron ZA, McClelland GB, and Scott GR (2019). Evolution of physiological  
808 performance capacities and environmental adaptation : insights from high-elevation  
809 deer mice ( *Peromyscus maniculatus* ). *J Mammal* 100:910–922.

810 Swenson ER, Maggiorini M, Mongovin S, Gibbs JSR, Greve I, Mairbäurl H, and  
811 Bartsch P (2002). Pathogenesis of High-Altitude Pulmonary Edema Inflammation is  
812 not an etiologic factor. *JAMA* 287.

813 Thomas SP, Lust MR, and Van Riper HJ (1984). Ventilation and Oxygen Extraction in  
814 the Bat *Phyllostomus hastatus* during Rest and Steady Flight. *Soc Integr Comp Biol*  
815 57.

816 Thomas SP, Follette DB, and Thomas GS (1995). Metabolic and ventilatory  
817 adjustments and tolerance of the bat *Pteropus poliocephalus* to acute hypoxic stress.  
818 *Comp Biochem Physiol A Physiol* 112:43–54.

- 819 Tucker VA (1968). Respiratory physiology of house sparrows in relation to high-  
820 altitude flight. *J Exp Biol* 48:55–66.
- 821 Tucker VA (1970). Energetic cost of locomotion in animals. *Comp Biochem Physiol*  
822 34:841–846.
- 823 Tucker A, McMurtry I, Reeves J, Alexander A, Will D, and Grover R (1975). Lung  
824 vascular smooth muscle as a determinant of pulmonary hypertension at high altitude.  
825 *Am J Physiol* 228:762–767.
- 826 Villafuerte FC and Corante N (2016). Chronic Mountain Sickness: Clinical Aspects,  
827 Etiology, Management, and Treatment. *High Alt Med Biol* 17:61–69.
- 828 Vogel JA and Harris CW (1967). Cardiopulmonary responses of resting man during  
829 early exposure to high altitude. *J Appl Physiol* 22:1124–1128.
- 830 Wagner PD (2010). Operation Everest II. *High Alt Med Biol* 11:111–119.
- 831 Ward S, Bishop CM, Woakes AJ, and Butler PJ (2002). Heart rate and the rate of  
832 oxygen consumption of flying and walking barnacle geese (*Branta leucopsis*) and bar-  
833 headed geese (*Anser indicus*). *J Exp Biol* 205:3347–3356.
- 834 Watson RR, Fu Z, and West JB (2007). Morphometry of the extremely thin pulmonary  
835 blood-gas barrier in the chicken lung. *Am J Physiol Lung Cell Mol Physiol* 292:L769–  
836 L777.
- 837 Watson RR, Fu Z, and West JB (2008). Minimal distensibility of pulmonary capillaries  
838 in avian lungs compared with mammalian lungs. *Respir Physiol Neurobiol* 160:208–  
839 214.
- 840 Weatheres W and Snyder G (1974). Functional Acclimation of Japanese Quail to  
841 Simulated High-Altitude. *J Comp Physiol* 137:127–137.

- 842 Weber JM, Roberts TJ, Vock R, Weibel ER, and Taylor CR (1996). Design of the  
843 oxygen and substrate pathways. III. Partitioning energy provision from carbohydrates.  
844 J Exp Biol 199:1659–1666.
- 845 Weber RE, Jessen T-H, Malte H, and Tame J (1985). Mutant hemoglobins (Ala-  $\alpha$  119  
846 and Pro- $\alpha$ 119 ): functions related to high-altitude respiration. J Appl Physiol 75:2646–  
847 2655.
- 848 Weinstein Y, Bernstein MH, Bickler PE, Gonzales DV, Samaniego FC, and Escobedo  
849 M. (1985). Blood respiratory properties in pigeons at high altitudes: effects of  
850 acclimation. Am J Physiol Integr Comp Physiol 249.
- 851 West JB (2002). Highest permanent human habitation. High Alt Med Biol 3:401–407.
- 852 West JB (2006). Human responses to extreme altitudes. Integr Comp Biol 46:25–34.
- 853 West JB (2009). Comparative physiology of the pulmonary blood-gas barrier: the  
854 unique avian solution. Am J Physiol Regul Integr Comp Physiol 297:R1625–R1634.
- 855 West JB, Watson RR, and Fu Z (2007). Major differences in the pulmonary circulation  
856 between birds and mammals. 157:382–390.
- 857 Williams TC, Ireland LC, and Williams JM (1973). High altitude flights of the free-tailed  
858 bat, *Tadarida brasiliensis*, observed with radar. J Mammal 54:807–821.
- 859 Willmer P, Stone G, and Johnston I (2005). Environmental Physiology of Animals.  
860 Blackwell publishing: Oxford.
- 861 Wilson MH and Imray CHE (2016). The cerebral venous system and hypoxia. J Appl  
862 Physiol 120:244–250.
- 863 Wolfel EE, Groves BM, Brooks GA, Butterfield GE, Mazzeo RS, Moore LG, Sutton JR,



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

864 Bender PR, Dahms TE, McCullough RE, McCullough RG, Huang S, Sun S, Grover  
865 RF, Hultgren H., and Reeves JT (1991). Oxygen transport during steady-state  
866 submaximal exercise in chronic hypoxia. J Appl Physiol 70:1129–1136.


867 Yap KN, Dick MF, Guglielmo CG, and Williams TD (2018). Effects of experimental  
868 manipulation of hematocrit on avian flight performance in high- and low-altitude  
869 conditions. J Exp Biol 221:jeb191056.

870 Yersin AAG, Huff WE, Kubena LF, Elissalde MH, Harvey RB, and Giroir LE (1992).  
871 Changes in Hematological, Blood Gas, and Serum Biochemical Variables in Broilers  
872 during Exposure to Simulated High. Avian Dis 36:189–196.

873 York JM, Chua BA, Ivy CM, Alza L, Cheek R, Scott GR, McCracken KG, Frappell PB,  
874 Dawson NJ, Laguë SL, and Milsom WK (2017). Respiratory mechanics of eleven avian  
875 species resident at high and low altitude. J Exp Biol 220:1079–1089.

876





Schematic diagram showing records of humans and birds found at high altitude in relation to commercial aircraft cruising altitude and some of the world's highest mountains. High-altitude birds are, for the majority of the time, found at much lower altitudes, but make rare flights above 5,500 metres altitude. Sources: Beall, 2014; Borrás et al., 2010; Dawson et al., 2016; Hawkes et al., 2011; Hawley et al., 2018; Laybourne, 1974; McCracken et al., 2009b; Mukherjee et al., 2008; Noel, 1927; Parr et al., 2017; West, 2002

174x102mm (150 x 150 DPI)




Diagram of the oxygen transport chain, detailing the physiological adjustments to high-altitude hypoxia that have been documented in low-altitude native humans ascending to high altitude and details of preadaptations that are common across all birds and allow for a greater tolerance of hypoxia relative to mammals before any physiological adjustments occur upon exposure to hypoxia. Sources: [1] West, 2006, [2] Moudgil et al., 2005, [3] Grocott et al., 2009, [4] Wagner, 2010, [5] Hoiland et al., 2018, [6] Maina, 2000, [7] Scott, 2011, [8] Scheid and Piiper, 1972, [9] Lague et al., 2016, [10] Grubb, 1983, [11] Maginniss et al., 1997, [12] Scott et al., 2009, [13] Faraci et al., 1984, [14] McGuire and Guglielmo, 2009, [15] Ward et al., 2002.

193x158mm (150 x 150 DPI)